



HYPERTENSION (HIGH BLOOD PRESSURE)

This sheet is for your information and is not a substitute for medical advice. You should contact your doctor or other healthcare provider with any questions about your health, treatment or care.

What is hypertension?

Hypertension is the medical term used to describe an elevated blood pressure, commonly referred to as *high blood pressure*. Everybody has and needs blood pressure. Without it, blood cannot circulate through the body, and without circulating blood, vital organs cannot obtain the oxygen and nutrients that they need to work. Normal blood pressure falls within a specific range – it is not restricted to one set of numbers.

When the heart beats, it pumps blood to the arteries and creates pressure in them. This pressure (blood pressure) is a result of two forces. The first force is created as blood pumps into the arteries and through the circulatory system. The second is created as the arteries resist the blood flow. If you are healthy, your arteries are muscular and elastic. They stretch when the heart pumps blood through them. How much they stretch depends on how much force the blood exerts.

Your heart beats about 60 to 80 times a minute under normal conditions. Your blood pressure rises with each heartbeat and falls when your heart relaxes between beats. Your blood pressure can change from minute to minute, with changes in posture, exercise or sleeping, but it should normally be less than 140/90mm Hg for an adult. Blood pressure that stays above this level is considered high. Your doctor may take several readings before making a judgment about whether your blood pressure is considered to be high.

What do blood pressure numbers indicate?

- The upper (systolic) number represents the pressure while the heart is pumping.
- The lower (diastolic) number represents the pressure when the heart is resting between beats.

The systolic pressure is always stated first and the diastolic pressure second. For example: 122/76 (122 over 76) means systolic = 122, diastolic = 76. The systolic is usually the higher number.

What should your blood pressure level be?

The table below provides a guideline for what is considered normal and high blood pressure levels:

Blood pressure category	Systolic (mm Hg)		Diastolic (mm Hg)	Follow-up recommended
Optimal	Less than 120	and	less than 80	Recheck in two years
Normal	Less than 130	and	less than 85	Recheck in two years
High normal	130 to 139	or	85 to 89	Recheck in one year

HIGH

Stage 1	140 to 159	or	90 to 99	Confirm within two months
Stage 2	160 to 179	or	100 to 109	Evaluate within one month
Stage 3	180 or higher	or	110 or higher	Evaluate immediately

IMPORTANT: Your doctor should evaluate unusually low readings.

What causes hypertension?

In 90-95% of high blood pressure cases, the cause is unknown. In fact, you can have high blood pressure for years without knowing it. That is why it's called 'the silent killer'. When the cause is unknown, you have what is called *essential* or *primary* hypertension.

You may be at risk of getting high blood pressure if you:

- are overweight
- eat too much salt
- regularly consume large quantities of alcohol
- do not exercise enough
- are constantly under stress
- have a family history of high blood pressure
- are 40 years or older
- have certain medical problems such as kidney diseases.

Some people have high blood pressure caused by underlying conditions which are known as *secondary* hypertension. Various conditions and medications can lead to secondary hypertension. This includes:

- kidney abnormalities
- a structural abnormality of the large blood vessel leaving the heart, called the aorta, existing since birth
- narrowing of arteries entering the kidney
- certain medications, such as birth control pills, cold remedies, decongestants, over-the-counter pain relievers and some prescription drugs
- illegal drugs, such as cocaine and amphetamines.

These problems can be corrected. For example, doctors can repair a narrowed artery that supplies blood to a kidney. Most of these problems can be ruled out by obtaining a thorough history, undergoing a physical examination and the taking of a few tests. Special tests are sometimes needed, but hospitalisation is not usually necessary.

How is blood pressure measured?

Blood pressure is measured using a medical instrument called a *sphygmomanometer*. A cuff is wrapped around a person's upper arm and inflated. When the cuff is inflated, it compresses a large artery in the arm, momentarily stopping the blood flow.

Next, air in the cuff is released and the person measuring the blood pressure listens with a stethoscope. When the blood starts to pulse through the artery, it makes a sound. Sounds continue to be heard until pressure in the artery exceeds the pressure in the cuff.

While the person listens and watches the gauge, he or she records two measurements. Systolic pressure is the pressure of the blood flow when the heart beats (the pressure when the first sound is heard). Diastolic pressure is the pressure between heartbeats (the pressure when the last sound is heard). Blood pressure is measured in millimetres of mercury, which is abbreviated mm Hg.

What is the effect of high blood pressure on your body?

High blood pressure adds to the workload of your heart and arteries. Your heart must pump harder and the arteries carry blood that is moving under greater pressure. For example, if you put too much pressure in a car tyre, it will inflate to a certain point and then it will burst. Similarly, if the arteries are functioning at extremely high pressures, they may rupture.

If high blood pressure continues for a long period of time, your heart and arteries may not function as well as they should be. Other body organs may also be affected. There is an increased risk of stroke, congestive heart failure, kidney failure and heart attack.

When high blood pressure exists with obesity, smoking, high blood cholesterol levels and/or diabetes, the risk of heart attack or stroke increases several times.

What are the symptoms of hypertension?

High blood pressure usually has no symptoms. In fact, many people have high blood pressure for years without knowing it. It doesn't refer to being tense, nervous or hyperactive. You can be a calm, relaxed person and still have high blood pressure. The only way to find out if you have high blood pressure is to have it checked by a healthcare practitioner. A blood pressure test is quick and painless. It can be done in a doctor's office, hospital clinic, school, nurse's office, company clinic or at a health and wellness day.

A single reading showing high blood pressure does not mean you have hypertension, but it is a sign that it should be monitored carefully. If your blood pressure is normal, have it checked at least every two years. If your blood pressure is near the top of the normal range, or if you have a family history of high blood pressure, you are at higher risk. Your doctor will tell you how often to have it checked.

What is the treatment for hypertension?

Lifestyle changes

Any of these can help lower high blood pressure:

- Have your blood pressure checked regularly.
- Maintain a healthy diet.
- Use less salt in your diet.
- Maintain a healthy weight – excess body weight is related to high blood pressure, and a weight loss programme is beneficial for lowering blood pressure and improving the effectiveness of medication for high blood pressure. Research has shown that weight reduction, even as little as 5kg, reduces high blood pressure in an overweight person.
- Be physically active for at least 30 minutes a day.
- Limit your alcohol intake.
- Stop smoking.

Drug treatment

Treatment is varied and changing all the time, with new drugs appearing on the market almost daily. There are several classes of drugs which act on different areas of the circulatory system and result in control of blood pressure. The choice of drug is affected by many factors, especially when you have other chronic conditions. It is important to work closely with your doctor to find the appropriate medication and treatment plan for you.

It is important to record your blood pressure and other relevant measurements and keep this information handy when seeing your doctor or when dealing with your medical scheme's chronic disease management programme:

- Blood pressure readings (e.g. 130/86)
- Your weight in kg
- Total cholesterol readings (mmol/l)
- Any emergency room visits
- Any doctor visits
- Additional notes and relevant measurements

References

1. AMERICAN SOCIETY OF HYPERTENSION. 2010. *Blood pressure and your health*. Website. <https://www.heart.org/en>.
2. CENTERS FOR DISEASE CONTROL AND PREVENTION (CDC). 22 October 2020. *High blood pressure*. Website. <https://www.cdc.gov/bloodpressure/index.htm>.
3. HEART AND STROKE FOUNDATION SOUTH AFRICA, THE. Website. <https://www.heartfoundation.co.za/>
4. SOUTHERN AFRICAN HYPERTENSION SOCIETY. Website. <https://www.hypertension.org.za/>.

The content of this publication is the intellectual property of Momentum Health (Pty) Limited and any reproduction of this publication is prohibited unless written consent is obtained.